

GAMMASPHERE STUDIES OF ^{252}Cf SPONTANEOUS FISSION. M.A. Stoyer, J.A. Becker, L.A.

Bernstein, R.W. Loughheed, K.J. Moody, and J.F. Wild, *Lawrence Livermore National Laboratory, Livermore, CA 94550*; J.O. Rasmussen, S.J. Asztalos, S.Y. Chu, R.B. Firestone, K.E. Gregorich, I.Y. Lee, A.O. Macchiavelli, and R.W. MacLeod, *Lawrence Berkeley National Laboratory, Berkeley CA 94720*; Y.X. Dardenne, R. Aryaeinejad, K. Butler-Moore, J.D. Cole, and M.W. Drigert, *Idaho National Engineering Laboratory, Idaho Falls, ID 83415*; R. Donangelo, *Universidade Federal do Rio de Janeiro, 21944 Rio de Janeiro*; J.H. Hamilton, B.R.S Babu, T.N. Ginter, J. Kormicki, and A.V. Ramayya, *Vanderbilt University, Nashville, TN 37235*; Yuri Oganessian, and Gurgen Ter-Akopian, *Joint Institute for Nuclear Studies, Dubna, Russia*; W.C. Ma, and P.G. Varrette, *Mississippi State University, Mississippi State, MS 39762*; and H.C. Griffin, *University of Michigan, Ann Arbor, MI 48109*. For the past 4 years, our group has been studying the spontaneous fission process by detecting gamma-rays emitted from fission fragments in various early implementations of Gammasphere. Both ^{242}Pu and ^{252}Cf have been studied. Some preliminary results will be discussed in this presentation, including for example zero-neutron fission, many-neutron fission and nuclear spectroscopy of neutron-rich nuclei. In the fall of 1996, ^{252}Cf will again be studied using the parallel-plate avalanche counters of the University of Rochester to detect the fission fragments in coincidence with gamma-rays, enabling direct detection of all fission products except for neutrons.

This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.